

Leading order corrections to the cosmological evolution of tensor perturbations in braneworld

Tsutomu Kobayashi* and Takahiro Tanaka†

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

Tensor type perturbations in the expanding brane world of the Randall Sundrum type are investigated. We consider a model composed of slow-roll inflation phase and the succeeding radiation phase. The effect of the presence of an extra dimension through the transition to the radiation phase is studied, giving an analytic formula for leading order corrections.

PACS numbers: 98.80.Cq, 04.50.+h

I. INTRODUCTION

In recent years there has been much interest in braneworld scenarios [1, 2, 3], where our four dimensional universe could be a brane embedded in a higher dimensional bulk spacetime. The basic ingredient of braneworld scenarios is that while ordinary matter fields are confined to the brane, only gravitational interactions can propagate in the bulk. The second Randall-Sundrum model (RS II) [4] with one brane in an anti de Sitter (AdS) bulk is particularly interesting in the point that, despite an infinite extra dimension, four dimensional general relativity (4D GR) can be recovered at low energies/long distances on the brane. Then, what is leading order corrections to the conventional gravitational theory? It was shown that the Newtonian potential in the RS II braneworld, including the correction due to the bulk gravitational effects with a precise numerical factor [5, 6], is given by

$$V(r) \simeq -\frac{Gmm'}{r} \left(1 + \frac{2\ell^2}{3r^2} \right).$$

Here ℓ is the curvature length of the AdS space and is experimentally constrained to be $\ell \lesssim 0.1$ mm. It is natural to ask next what are leading order corrections to cosmological perturbations.

Great efforts have been paid for the problem of calculating cosmological perturbations in braneworld scenarios (see, for example, Refs. [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27]). In order to correctly evaluate perturbations on the brane, we need to solve bulk perturbations, which reduces to a problem of solving partial differential equations with appropriate boundary conditions. Hence it is not as easy as in the standard four dimensional cosmology. A pure de Sitter braneworld [15], and its variations such as “junction” models [16, 17] and a special dilatonic braneworld [24, 25, 26] are the rare examples where the bulk perturbations are analytically solved. In more generic cases, one has to resort to a numerical calculation [18, 21, 22], or some approximate methods [14, 19, 20, 23]. There is another difficulty concerned with a physical, fundamental aspect of the problem; we do not know how to specify appropriate initial conditions for perturbations with bulk degrees of freedom.

In this paper, we investigate cosmological tensor perturbations in the RS II braneworld, and try to evaluate leading order corrections to the 4D GR result analytically. For this purpose, we make use of the reduction scheme to a four dimensional effective equation which iteratively takes into account the effects of the bulk gravitational fields [28]. As a first step, we concentrate on perturbations which is initially at a super-horizon scale.

The paper is organized as follows. In Sec. II we define the setup of the problem discussed in the present paper more precisely. In Sec. III we briefly review the derivation of the effective equation of motion for tensor perturbations. In Sec. IV, using the long wavelength approximation we develop a method for evaluating leading order corrections, and evaluate the corrections considering slow-roll inflation and the radiation stage which follows after inflation. The shortcoming of the long wavelength approximation is cured by introducing another method suitable for analyzing the radiation phase in Sec. IV. Section V is devoted to summary and discussion.

*Electronic address: tsutomu@tap.scphys.kyoto-u.ac.jp

†Electronic address: tama@scphys.kyoto-u.ac.jp

II. SETUP OF THE PROBLEM AND BACKGROUND COSMOLOGICAL MODEL

In this paper we investigate tensor perturbations in a Friedmann braneworld model with an warped (infinite) extra dimension [4, 29, 30, 31, 32, 33]. To determine the spectrum of tensor perturbations observed at late time, we need to solve the perturbation equations with appropriate initial conditions. However, it is not an easy task to specify what initial conditions we should use in the context of braneworld cosmology. At an earlier stage of the evolution of the universe the physical wavelength of perturbation modes is much shorter than the bulk curvature scale. Then, the correction to the four dimensional standard evolution will not remain small. Therefore, in this regime it is not appropriate to discuss the dynamics of perturbation modes under the assumption that a modification to the four dimensional standard one is small. Hence our strategy is to isolate the issue of initial conditions from the rest. In this paper we discuss a relatively easy part, *i.e.*, late time evolution after the effect of the fifth dimension becomes perturbative.

Our discussion will be restricted to the perturbation modes whose wavelength is initially much longer than the Hubble horizon scale. We analytically derive a formula for small corrections to the solution for the tensor perturbation equation due to the presence of the fifth dimension. We mainly focus on the behavior of a growing solution, since a growing solution will be more important than a decaying one in general. Later we briefly mention the decaying solution in Sec. VI.

The background spacetime that we consider is composed of a five dimensional AdS bulk, whose metric is given in Poincaré coordinates as

$$ds^2 = \frac{\ell^2}{z^2} (dz^2 - dt^2 + \delta_{ij} dx^i dx^j),$$

with a Friedmann brane at $z = z(t)$. Here ℓ is the curvature length of AdS space. In the original RS II model [4], a brane is placed at a position of fixed z , and Minkowski geometry is realized on the brane. The induced metric on $z = z(t)$ is

$$\begin{aligned} ds^2 &= a^2(t) [-(1 - \dot{z}^2) dt^2 + \delta_{ij} dx^i dx^j], \\ a(t) &= \ell/z(t), \end{aligned} \quad (1)$$

where the overdot denotes ∂_t . From Eq. (1), we see that the conformal time on the brane is given by $d\eta = \sqrt{1 - \dot{z}^2} dt$. The brane motion is related to the energy density of matter localized on the brane by the modified Friedmann equation [29, 30, 31, 32, 33] as

$$H^2 = \frac{8\pi G}{3} \left(\rho + \frac{\rho^2}{2\sigma} \right), \quad (2)$$

and

$$H := \frac{\dot{a}}{a} = \frac{-\ell^{-1} \dot{z}}{\sqrt{1 - \dot{z}^2}}, \quad (3)$$

where $\sigma = 3/4\pi G\ell^2$ is a tension of the brane. The quadratic term in ρ in the Friedmann equation modifies the standard cosmological expansion law. At low energies ($\ell H \ll 1$), the brane motion is non-relativistic ($\dot{z} \ll 1$), and hence the conformal time on the brane η almost coincides with the bulk conformal time t .

As a background Friedmann brane model, we consider slow-roll inflation at low energies followed by a radiation dominant phase. During the slow-roll inflation the wavelength of perturbations stays outside the Hubble scale, and it re-enters the Hubble horizon only during the radiation era.

We fix a model of slow-roll inflation by simply providing a function

$$\epsilon(\eta) := 1 - \frac{\partial_\eta \mathcal{H}}{\mathcal{H}^2}, \quad (4)$$

where $\mathcal{H} := \partial_\eta a/a = aH$. During slow-roll inflation, ϵ is supposed to be small. In the $\epsilon \rightarrow 0$ limit, we recover the de Sitter case

$$a = \frac{a_0 \eta_0}{2\eta_0 - \eta}, \quad \mathcal{H} = \frac{1}{2\eta_0 - \eta}, \quad (\eta < \eta_0), \quad (5)$$

where we have chosen the normalization of the scale factor and the origin of the η -coordinate so that $a = a_0$ and $\mathcal{H} = 1/\eta_0$ at $\eta = \eta_0$.

In the above alternatively we could specify the potential of the inflaton field localized on the brane. Then, for a given inflaton potential, the evolution of the background scale factor would differ from the conventional four dimensional one. However, as we will show below, the correction to the tensor perturbations coming from the slow-roll inflation phase is always small irrespective of the details of the evolution of the scale factor. Hence, we simply give time-dependence of the slow-roll parameter to specify a model.

We assume that slow-roll inflation terminates at around $\eta = \eta_0$, and a radiation era follows. During the radiation era the energy density decreases like $\rho = \rho_0(a_0/a)^4$ as usual. Then, solving Eq. (2) at low energies, we find that the scale factor behaves like

$$a(\eta) = {}^{(0)}a(\eta) + \delta a(\eta) + \mathcal{O}(\ell^4), \quad (\eta > \eta_0), \quad (6)$$

with

$${}^{(0)}a(\eta) = a_0 \frac{\eta}{\eta_0}, \quad \delta a(\eta) = a_0 \ell^2 H_0^2 \left[\frac{1}{6} - \frac{1}{24} \left(\frac{\eta_0}{\eta} \right)^3 - \frac{1}{8} \left(\frac{\eta}{\eta_0} \right) \right], \quad (7)$$

and $H_0 = (a_0 \eta_0)^{-1}$. Correspondingly, we have

$$\mathcal{H}(\eta) = {}^{(0)}\mathcal{H}(\eta) + \delta \mathcal{H}(\eta) + \mathcal{O}(\ell^4), \quad (8)$$

with

$${}^{(0)}\mathcal{H}(\eta) = \frac{1}{\eta}, \quad \delta \mathcal{H}(\eta) = \frac{\ell^2}{6a^2 \eta_0^3} \left[\left(\frac{\eta_0}{\eta} \right)^3 - 1 \right], \quad (\eta > \eta_0). \quad (9)$$

It is convenient to extend the definition of the slow-roll parameter ϵ to the later epoch by

$$\epsilon(\eta) := 1 - \frac{\partial_\eta {}^{(0)}\mathcal{H}}{{}^{(0)}\mathcal{H}^2}. \quad (10)$$

III. TENSOR PERTURBATIONS ON A FRIEDMANN BRANE

We use the method to reduce the five dimensional equation for tensor perturbations in a Friedmann braneworld at low energies to a four dimensional effective equation of motion, derived in Ref. [28]. Here we summarize the derivation.

Tensor perturbations on a Friedmann brane [15, 16, 17, 18, 19, 20, 21, 22, 23] are given by

$$ds^2 = \frac{\ell^2}{z^2} [dz^2 - dt^2 + (\delta_{ij} + h_{ij})dx^i dx^j]. \quad (11)$$

We expand the perturbations by using $Y_{ij}^k(\mathbf{x})$, a transverse traceless tensor harmonics with comoving wave number k , as $h_{ij} = \sum_k Y_{ij}^k(\mathbf{x}) \Phi_k(t, z)$. Then the equation of motion for the tensor perturbations in the bulk is given by

$$\left(-\partial_z^2 + \frac{3}{z} \partial_z + \partial_t^2 + k^2 \right) \Phi_k = 0, \quad (12)$$

Hereafter we will discuss each Fourier mode separately, and we will abbreviate the subscript k . The general solution of this equation is

$$\begin{aligned} \Phi &= \int d\omega \tilde{\Psi}(\omega) e^{-i\omega t} 2(pz)^2 K_2(pz) \\ &= \int d\omega \tilde{\Psi}(\omega) e^{-i\omega t} \left\{ 1 - \frac{(pz)^2}{4} + \frac{(pz)^4}{16} \left[\frac{3}{4} - \gamma - \ln \left(\frac{pz}{2} \right) \right] + \dots \right\}, \end{aligned} \quad (13)$$

where $p^2 = -\omega^2 + k^2$, and γ is Euler's constant. We have chosen the branch cut of the modified Bessel function K_2 so that there is no incoming wave from past null infinity in the bulk. The coefficients $\tilde{\Psi}(\omega)$ are to be determined by the boundary condition on the brane $n^\mu \partial_\mu \Phi|_{z=z(t)} = 0$, where n^μ is a unit normal to the brane, or, equivalently, by the effective Einstein equations on the brane [34],

$${}^{(4)}G_{\mu\nu} = 8\pi G T_{\mu\nu} + (8\pi G_5)^2 \pi_{\mu\nu} - E_{\mu\nu}. \quad (14)$$

Here $\pi_{\mu\nu}$ is quadratic in the energy momentum tensor $T_{\mu\nu}$, and the first two terms on the right hand side are totally represented by the variables which reside on the brane. A projected Weyl tensor $E_{\mu\nu} := C_{\alpha\mu\beta\nu}n^\alpha n^\beta$ represents the effects of the bulk gravitational fields, giving rise to corrections to four dimensional Einstein gravity in a fairly nontrivial way. In the present case, this can be written explicitly, and the effective four dimensional equation reduces to

$$(\partial_\eta^2 + 2\mathcal{H}\partial_\eta + k^2)\phi = -2E, \quad (15)$$

$$-2E = \left[(\ell H)^2 (\partial_t^2 + \partial_z^2) - 2\ell H \sqrt{1 + (\ell H)^2} \partial_t \partial_z + \partial_z^2 - \frac{1}{z} \partial_z \right] \Phi \Big|_{z=z(t)}, \quad (16)$$

where $\phi(t) := \Phi(t, z(t))$ is the perturbation evaluated on the brane. At low energies ($\ell H \ll 1$), we can use the approximation like

$$\partial_t^2 \Phi \simeq \partial_\eta^2 \Phi, \quad \partial_t \partial_z \Phi \simeq -\frac{\ell}{2a} \partial_\eta (\partial_\eta^2 + k^2) \Phi, \quad \partial_z^2 \Phi \simeq -\frac{1}{2} \int d\omega \tilde{\Psi} e^{-i\omega\eta} p^2 \simeq -\frac{1}{2} (\partial_\eta^2 + k^2) \Phi,$$

neglecting higher order corrections of $\mathcal{O}(\ell^4)$. Then, using the lower order equation to eliminate the higher derivative terms in $-2E$, we finally obtain our basic equation:

$$\left(\partial_\eta^2 + 2\overset{(0)}{\mathcal{H}}\partial_\eta + k^2 \right) \phi \simeq \frac{\ell^2}{a^2} (\mathcal{S}_0[\phi] + \mathcal{S}_1[\phi] + \mathcal{S}_2[\phi]), \quad (17)$$

where

$$\mathcal{S}_0[\phi] = -\frac{2\overset{(0)}{a}^2}{\ell^2} \delta\mathcal{H}\phi', \quad (18)$$

$$\mathcal{S}_1[\phi] = (3\overset{(0)}{\mathcal{H}}^3 - 2\overset{(0)}{\mathcal{H}}\overset{(0)}{\mathcal{H}}')\phi' + k^2\overset{(0)}{\mathcal{H}}^2\phi, \quad (19)$$

$$\mathcal{S}_2[\phi] = -\frac{1}{2} \int d\omega p^4 \tilde{\phi} e^{-i\omega\eta} \left[\ln\left(\frac{p\ell}{2a}\right) + \gamma \right], \quad (20)$$

and the prime denotes ∂_η . The first term arises due to the non-standard cosmological expansion included in $\mathcal{H}$, while the last two terms are the corrections from the bulk effects $E_{\mu\nu}$. We can see that all terms are suppressed by ℓ^2 (or $\ell^2 \ln \ell$)¹. $\mathcal{S}_2$ is essentially nonlocal because of the presence of the log term. A time-domain expression for the quantity $p^4\phi$ which appears in $\mathcal{S}_2$ can be rewritten, by eliminating the higher derivative terms in a similar way as before, as

$$\hat{p}^4\phi \simeq -2 \left[\left(\overset{(0)}{\mathcal{H}}'' - 6\overset{(0)}{\mathcal{H}}\overset{(0)}{\mathcal{H}}' + 4\overset{(0)}{\mathcal{H}}^3 \right) \partial_\eta + 2k^2 \left(\overset{(0)}{\mathcal{H}}^2 - \overset{(0)}{\mathcal{H}}' \right) \right] \phi, \quad (21)$$

where we have introduced a derivative operator $\hat{p}^2 = \partial_\eta^2 + k^2$.

IV. NEW LONG WAVELENGTH APPROXIMATION

Let us begin with the case that the wavelength of perturbations is much longer than the Hubble scale. The long wavelength approximation not only allows us an easy treatment of perturbations, but also turns out to be sufficient to derive all the major corrections except for that coming from $\mathcal{S}_0$. We would like to stress that what we call the long wavelength approximation here is different from a low energy expansion scheme or a gradient expansion scheme in the literature [35, 36, 37].

A. General iteration scheme

The equation we are considering takes the form of

$$\left(\partial_\eta^2 + 2\overset{(0)}{\mathcal{H}}\partial_\eta + k^2 \right) \phi = \frac{\ell^2}{a^2} \mathcal{S}[\phi]. \quad (22)$$

¹ Hereafter, we will refer to both terms suppressed by ℓ^2 and by $\ell^2 \ln \ell$ as $\mathcal{O}(\ell^2)$.

This can be solved iteratively taking ℓ^2 as a small parameter. For this purpose, we write

$$\phi(\eta) = \phi^{(0)}(\eta) e^{F(\eta)} = \phi^{(0)}(\eta) \exp \left[\int^\eta f(\eta') d\eta' \right]. \quad (23)$$

A zeroth order solution $\phi^{(0)}$ by definition satisfies the equation obtained by setting the right hand side of Eq. (22) to be zero. Then f obeys

$$\partial_\eta f + 2 \left(\frac{\partial_\eta \phi^{(0)}}{\phi^{(0)}} + \mathcal{H} \right) f = \frac{\ell^2}{a^{(0)2} \phi^{(0)}} \mathcal{S}[\phi^{(0)}], \quad (24)$$

which can be integrated immediately to give

$$f(\eta) = \frac{\ell^2}{a^{(0)2} \phi^{(0)}} \int^\eta d\eta' \phi^{(0)} \mathcal{S}[\phi^{(0)}]. \quad (25)$$

Integrating this expression, we obtain the first order correction F . In the present case, $\mathcal{S}$ consists of three parts, $\mathcal{S}_0$, $\mathcal{S}_1$ and $\mathcal{S}_2$. We hereafter denote the corrections f and F coming from $\mathcal{S}_i$ as f_i and F_i , respectively.

B. Long wavelength expansion of the zeroth order solution

In the long wavelength approximation, we can write the zeroth order solution explicitly as

$$\phi^{(0)} \simeq A_k \left[1 - k^2 \int^\eta d\eta' I(\eta') \right], \quad (26)$$

$$\partial_\eta \phi^{(0)} \simeq -A_k k^2 I(\eta), \quad (27)$$

where A_k is an amplitude of the perturbation which can depend on k , and

$$I(\eta) := a^{(0)-2}(\eta) \int_{-\infty}^\eta a^{(0)2}(\eta') d\eta'. \quad (28)$$

Here we simply keep the terms up to $\mathcal{O}(k^2)$.

We have already introduced a parameter $\epsilon(\eta)$ in Eq. (10), which parametrizes the cosmological expansion. In the inflationary universe, it reduces to a slow-roll parameter and is assumed to be small. Slow-roll inflation assumes that time differentiation of ϵ is $\mathcal{O}(\epsilon^2)$. Hence, we treat ϵ as a constant during slow-roll inflation. The transition from inflation to the radiation stage occurs at around $\eta = \eta_0$, and the behavior of ϵ there depends on the details of the reheating process. During the radiation stage, ϵ keeps again a constant value, $\epsilon = 2$, because the scale factor behaves like $a^{(0)} \propto \eta$. In general ϵ reduces to a constant for the scale factor $a^{(0)}$ proportional to a power of the conformal time.

Using ϵ and the zeroth order solution in the long wavelength approximation, we can write $\mathcal{S}_1[\phi^{(0)}]$ and $\hat{p}^4 \phi^{(0)}$ as

$$\begin{aligned} \mathcal{S}_1[\phi^{(0)}] &= \mathcal{H}^2 \left[(1 + 2\epsilon) \mathcal{H} \partial_\eta \phi^{(0)} + k^2 \phi^{(0)} \right] \\ &\simeq A_k k^2 \mathcal{H}^2 \left[1 - \mathcal{H} I - 2\epsilon \mathcal{H} I \right], \end{aligned} \quad (29)$$

$$\begin{aligned} \hat{p}^4 \phi^{(0)} &\simeq 2 \mathcal{H}^2 \left\{ \left[\epsilon' - 2\epsilon(1 + \epsilon) \mathcal{H} \right] \partial_\eta \phi^{(0)} - 2\epsilon k^2 \phi^{(0)} \right\} \\ &\simeq -2A_k k^2 \mathcal{H}^2 \left\{ \epsilon' I - 2\epsilon \left[(1 + \epsilon) \mathcal{H} I - 1 \right] \right\}, \end{aligned} \quad (30)$$

which are generic expressions not relying on any specific time dependence of the scale factor.

Equation (30) can be further rewritten into a more suggestive form. First, integrating the identity $\partial_\eta (a^{(0)2} / \mathcal{H}) = (1 + \epsilon) a^{(0)2}$, it can be shown that

$$1 - \mathcal{H} I = \frac{\mathcal{H}}{a^{(0)2}} \int_{-\infty}^\eta \epsilon a^{(0)2} d\eta'.$$

Then, integration by parts leads

$$(1 + \epsilon)^{(0)} \mathcal{H} I - 1 = \frac{\mathcal{H}}{a^{(0)2}} \int_{-\infty}^{\eta} \epsilon' a^{(0)2} I d\eta', \quad (31)$$

and thus we have

$$\hat{p}^4 \phi^{(0)} \simeq -2A_k k^2 \mathcal{H}^2 \left(\epsilon' I - 2\epsilon \frac{\mathcal{H}}{a^{(0)2}} \int_{-\infty}^{\eta} \epsilon' a^{(0)2} I d\eta' \right). \quad (32)$$

Since ϵ' is of second order in the slow-roll parameters during slow-roll inflation, we can safely neglect this term until the end of inflation. Moreover, for sufficiently smooth ϵ this term is expected to be small, and $\epsilon' = 0$ for the scale factor proportional to a power of the conformal time. Only at the time when the equation of state of the universe changes abruptly, *i.e.*, at a sudden transition from inflation to the radiation stage, ϵ' possibly becomes significantly large like a delta function. After such a violent transition, the integral in Eq. (32) leaves a constant value and thus this second term decreases in proportion to the damping factor $\mathcal{H}/a^{(0)2}$.

C. Slow-roll inflation

Now let us consider slow-roll inflation on the brane. In this case $\mathcal{S}_0$ vanishes, since $\delta\mathcal{H} = 0$ by construction. In the previous section we have also shown that $\hat{p}^4 \phi^{(0)}$ vanishes at first order in the slow-roll parameters during inflation. Consequently, the only relevant term in the present situation is $\mathcal{S}_1$.

Equation (29) together with Eq. (31) tells us that

$$\mathcal{S}_1 \simeq -A_k k^2 \mathcal{H}^2 \epsilon, \quad (33)$$

and thus

$$f(\eta) \simeq -\frac{k^2 \ell^2}{a^{(0)2}} \int_{-\infty}^{\eta} d\eta' \epsilon' \mathcal{H}^2. \quad (34)$$

Using the relations $\mathcal{H}^{(0)} \simeq a/a_0 \eta_0$ and $d\eta = a_0 \eta_0 d a^{(0)}/a^2$, and neglecting the time variation of the slow-roll parameter, we obtain

$$f(\eta) \simeq \frac{k^2 \ell^2}{\eta_0 a^{(0)2}} \epsilon \left(\frac{a_i}{a_0} - \frac{a}{a_0} \right), \quad (35)$$

where $a_i = a(\eta_i)$ and η_i is the lower boundary of integration. Integrating this expression, we have

$$F(\eta) \simeq -\frac{k^2 \ell^2}{a_i^2} \epsilon \left[\frac{1}{3} \left(\frac{a_i}{a^{(0)}} \right)^3 - \frac{1}{2} \left(\frac{a_i}{a^{(0)}} \right)^2 \right]. \quad (36)$$

We chose the integration constant so that F vanishes in the limit of $a(\eta) \rightarrow \infty$. This means that we have renormalized A_k so that it represents the amplitude of fluctuations that we see at late times if slow-roll inflation lasted forever.

Although our present long wavelength approximation is not valid when the wavelength $a_i k^{-1}$ becomes comparable or shorter than the Hubble scale H , we can still arrange a_i to be sufficiently small so that we can neglect the first term in Eq. (36). For such a_i , we have an expression independent of a_i ,

$$F(\eta) \approx \frac{k^2 \ell^2}{2 a^{(0)2}} \epsilon, \quad (37)$$

and we do not need to care about the choice of the “initial time” η_i . We see that the correction arising during inflation is very tiny, suppressed by the slow-roll parameter ϵ in addition to the factor $k^2 \ell^2 / a^{(0)2}$. For pure de Sitter inflation there is no correction, as is expected.




FIG. 1: Behavior of $\mathcal{S}_1$ around the transition time with the vertical axis in an arbitrary unit. Red solid line shows the case of a sharp transition with $s = 0.02\eta_0$, while blue dashed line represents the case of a smooth transition with $s = 0.5\eta_0$. The wavelength of the mode is chosen to be $k\eta_0 = 0.01$.




FIG. 2: Behavior of $\mathcal{S}_2$ around the transition time with the vertical axis in an arbitrary unit. Red solid line shows the case of a sharp transition with $s = 0.02\eta_0$, while blue dashed line represents the case of a smooth transition with $s = 0.5\eta_0$. The choice of the parameter is the same as in Fig. 1, $k\eta_0 = 0.01$.

D. Transition to the radiation stage

In this section, we investigate the effects of the transition from the inflation stage to the radiation stage. First, just for the illustrative purpose, we plot the behavior of $\mathcal{S}_1$ and $\mathcal{S}_2$ for the modes well outside the horizon in the neighborhood of the transition time η_0 in Fig. 1 and Fig 2. These plots are for ϵ given by

$$\epsilon(\eta) = \tanh[(\eta - \eta_0)/s] + 1, \quad (38)$$

where the parameter s controls the smoothness of the transition. From the figures it can be seen that the corrections from $E_{\mu\nu}$, $\mathcal{S}_1$ and $\mathcal{S}_2$, become significant only around the transition time.

From now on we shall consider the limiting case where ϵ is given by $\epsilon(\eta) = 2\theta(\eta - \eta_0)$. Here we neglected the tiny effect of the non vanishing slow-roll parameter during inflation. For this instantaneous transition model, the scale factor and the comoving Hubble parameter for $\eta < \eta_0$ are given by Eq. (5), while for $\eta > \eta_0$ they are given by $\overset{(0)}{a}$ and $\overset{(0)}{\mathcal{H}}$ in Eqs. (7) and (9).

Assuming such a step function-like transition, we can evaluate the corrections from $\mathcal{S}_0$ at $\eta > \eta_0$ as

$$f_0(\eta) \simeq -\frac{k^2 \ell^2}{9 a_0^2} \int_{\eta_0}^{\eta} \frac{\eta' d\eta'}{\eta_0^3} \left(1 - \frac{\eta_0^3}{\eta'^3}\right) \left(1 + 2 \frac{\eta_0^3}{\eta'^3}\right), \quad (39)$$

$$F_0(\eta) \simeq -\frac{k^2 \ell^2}{90 a_0^2} \left[5 \frac{\eta}{\eta_0} - 9 + 5 \left(\frac{\eta_0}{\eta}\right)^2 - \left(\frac{\eta_0}{\eta}\right)^5 \right]. \quad (40)$$

Here the integration constant is chosen so that the correction vanishes before the transition. From this expression, we find that F_0 diverges for a large η , and hence the correction to the final amplitude of fluctuations from $\mathcal{S}_0$ looks infinitely large. However, this is an artifact of the long wavelength approximation. At a late time during the radiation era the zeroth order solution becomes oscillatory. This oscillation suppresses the late time contribution. However, in the long wavelength approximation, this oscillation is not taken into account. In the succeeding section, we will develop another method which takes into account this oscillation. There we will find a finite correction.

As for $\mathcal{S}_1$, we have

$$f_1(\eta) \simeq -\frac{2}{3} \frac{k^2 \ell^2}{a_0^2} \int_{\eta_0}^{\eta} \frac{d\eta'}{\eta'^2} \left(1 + 5 \frac{\eta_0^3}{\eta'^3}\right), \quad (41)$$

$$F_1(\eta) \simeq -\frac{2}{3} \frac{k^2 \ell^2}{a_0^2} \left[\frac{3}{2} - \frac{9}{4} \left(\frac{\eta_0}{\eta}\right) + \frac{1}{2} \left(\frac{\eta_0}{\eta}\right)^2 + \frac{1}{4} \left(\frac{\eta_0}{\eta}\right)^5 \right]. \quad (42)$$

Long time after the transition (but of course before horizon re-entry), the correction becomes time independent,

$$\lim_{\eta \rightarrow \infty} F_1(\eta) = -\frac{k^2 \ell^2}{a_0^2}. \quad (43)$$

The next step is to evaluate the correction coming from $\mathcal{S}_2$. This can be done as follows. First, for the instantaneous transition $\hat{p}^4 \phi^{(0)}$ becomes

$$\hat{p}^4 \phi^{(0)} \simeq -4A_k k^2 \left[\frac{1}{\eta_0} \delta(\eta - \eta_0) - 4\theta(\eta - \eta_0) \frac{\eta_0^3}{\eta^5} \right] = -4A_k k^2 \partial_\eta \left[\theta(\eta - \eta_0) \frac{\eta_0^3}{\eta^4} \right]. \quad (44)$$

Then, integrating by parts, the Fourier transform of $\hat{p}^4 \phi^{(0)}$ is obtained as

$$\begin{aligned} p^4 \tilde{\phi}^{(0)} &= \frac{2A_k k^2}{\pi} i\omega \int d\eta' \theta(\eta - \eta_0) \frac{\eta_0^3}{\eta'^4} e^{i\omega\eta'} \\ &= \frac{2A_k k^2}{\pi} i\omega e^{i\omega\eta_0} q(\omega\eta_0), \end{aligned} \quad (45)$$

with

$$q(\omega\eta_0) := \int_1^\infty dy \frac{e^{i\omega\eta_0(y-1)}}{y^4}. \quad (46)$$

We separate $\mathcal{S}_2$ into two parts,

$$\mathcal{S}_2 = -\frac{1}{4} \int d\omega p^4 \tilde{\phi}^{(0)} e^{-i\omega\eta} \ln\left(\frac{p^2}{k^2}\right) - \frac{1}{2} \left[\gamma + \ln\left(\frac{k\ell}{2a_0}\right) + \ln\left(\frac{\eta_0}{\eta}\right) \right] \hat{p}^4 \phi^{(0)},$$

and, by substituting the expressions obtained above, we have

$$\mathcal{S}_2 = \frac{A_k k^2}{2\pi} \int d\omega q(\omega\eta_0) \partial_\eta \left[e^{-i\omega T} \ln\left(\frac{p^2}{k^2}\right) \right] + 2A_k k^2 \left[\gamma + \ln\left(\frac{k\ell}{2a_0}\right) + \ln\left(\frac{\eta_0}{\eta}\right) \right] \partial_\eta \left[\theta(T) \frac{\eta_0^3}{\eta^4} \right], \quad (47)$$

where $T := \eta - \eta_0$. Then, it can be integrated to give

$$\begin{aligned} f_2(\eta) &= \frac{k^2 \ell^2}{a_0^2} \left\{ \frac{1}{2\pi} \int d\omega q(\omega\eta_0) \left[e^{-i\omega T} \ln\left(\frac{p^2}{k^2}\right) \right] \right. \\ &\quad \left. + 2 \left[\gamma + \ln\left(\frac{k\ell}{2a_0}\right) + \ln\left(\frac{\eta_0}{\eta}\right) \right] \theta(T) \frac{\eta_0^3}{\eta^4} + \int_{-\infty}^{\eta} d\eta' \theta(\eta' - \eta_0) \frac{2}{\eta'} \frac{\eta_0^3}{\eta'^4} \right\}. \end{aligned} \quad (48)$$

The brunch cuts of log function and the path of ω integration in the first term should be chosen so that the retarded boundary condition is ensured, which are presented in Fig. 3. The integration is dominated by $\omega^2 \approx k^2$, where we may approximate $q(\omega\eta_0) \approx q(k\eta_0) \approx 1/3$ because $k\eta_0$ is assumed to be small. Then, the first term can be written by using the formula presented in Appendix, and can be integrated to give a part of $F_2(\eta)$ as

$$\begin{aligned} & -\frac{2k^2\ell^2}{3a_0^2} \int_{-\infty}^{\eta} d\eta' \frac{\eta_0^2}{\eta'^2} \left\{ \theta(\eta' - \eta_0) \frac{\cos[k(\eta' - \eta_0)] - 1}{\eta' - \eta_0} + \partial_{\eta'} [\theta(\eta' - \eta_0) \ln(k(\eta' - \eta_0))] + \gamma \delta(\eta' - \eta_0) \right\} \\ & \simeq -\frac{2k^2\ell^2}{3a_0^2} \left\{ \frac{\eta_0^2}{\eta^2} \ln(k(\eta - \eta_0)) + \ln(k\eta_0) \left(1 - \frac{\eta_0^2}{\eta^2} \right) - 1 + \frac{\eta_0}{\eta} + \ln \left(\frac{\eta - \eta_0}{\eta} \right) - \frac{\ln(\eta/\eta_0 - 1)}{(\eta/\eta_0)^2} + \gamma \right\} \\ & \xrightarrow{\eta \rightarrow \infty} -\frac{2k^2\ell^2}{3a_0^2} [\gamma - 1 + \ln(k\eta_0)], \end{aligned}$$

where we have neglected the term proportional to $\cos[k(\eta' - \eta_0)] - 1$ in the integrand because this term brings the higher order contributions in k . The other part of $F_2(\eta)$ is obtained by integrating the remaining parts in Eq. (48) as

$$\begin{aligned} & \frac{2k^2\ell^2}{a_0^2} \left\{ \left[\gamma + \ln \left(\frac{k\ell}{2a_0} \right) \right] \frac{1}{5} \left(1 - \frac{\eta_0^5}{\eta^5} \right) + \frac{1}{25} \left[\frac{\eta_0^5}{\eta^5} - 1 + 5 \frac{\ln(\eta/\eta_0)}{(\eta/\eta_0)^5} \right] + \frac{1}{4} \left(1 - \frac{\eta_0}{\eta} \right) - \frac{1}{20} \left(1 - \frac{\eta_0^5}{\eta^5} \right) \right\} \\ & \xrightarrow{\eta \rightarrow \infty} \frac{2k^2\ell^2}{a_0^2} \left\{ \frac{1}{5} \left[\gamma + \ln \left(\frac{k\ell}{2a_0} \right) \right] + \frac{4}{25} \right\}. \end{aligned}$$

Combining the above two, we finally obtain the correction from $\mathcal{S}_2$,

$$\lim_{\eta \rightarrow \infty} F_2(\eta) = \frac{2k^2\ell^2}{a_0^2} \left\{ \frac{37}{75} - \frac{2}{15} \left[\gamma + \ln \left(\frac{k}{a_0 H_0} \right) \right] + \frac{1}{5} \ln \left(\frac{\ell H_0}{2} \right) \right\}. \quad (49)$$

V. CORRECTIONS IN THE RADIATION STAGE

So far we have worked in the long wavelength approximation, and found k -dependent small corrections to the spectrum of tensor perturbations due to $\mathcal{S}_1$ and $\mathcal{S}_2$, which correspond to $E_{\mu\nu}$ term. It might be possible, however, that further corrections arise after the mode re-enters the horizon. We will show that such corrections are highly suppressed. A basic observation supporting this conclusion is that the contributions from $E_{\mu\nu}$ term in the right hand side of Eq. (17) become significantly large only at the transition time. It decreases in powers of $a(\eta)$ after the transition, and will be negligible when the long wavelength approximation brakes down.

In the long wavelength approximation, we could not obtain a meaningful estimate for the leading order correction caused by unconventional cosmic expansion, $\mathcal{S}_0$. Here we will give a more rigorous treatment, and will resolve this drawback of the approach taken in the preceding section.

In the radiation stage,

$$u(\eta) := a(\eta)\phi(\eta), \quad (50)$$




FIG. 3: Branch cuts and the contour of the integration in Eq. (48) for $T > 0$. For $T < 0$ we close the contour on the upper half plane.

is a convenient variable. In terms of this new variable, we can rewrite the equation of motion (17) as

$$u'' + k^2 u = \frac{\ell^2}{a} (\bar{\mathcal{S}}_0 + \mathcal{S}_1 + \mathcal{S}_2), \quad (51)$$

with

$$\bar{\mathcal{S}}_0 = \frac{a''}{\ell^2} u. \quad (52)$$

If we neglect the deviation of the expansion law from the standard one, we have $a \propto \eta$ in the radiation stage and so $a'' = 0$. This means that the first term on the right hand side gives a contribution of $\mathcal{O}(\ell^2)$ from the modification of the expansion law as $\bar{\mathcal{S}}_0$ in the preceding section. Thus, all the terms collected on the right hand side are of $\mathcal{O}(\ell^2)$, and at zeroth order u behaves just like a harmonic oscillator. The zeroth order solution is expressed by a linear combination of the standard modes:

$$u^{(0)}(\eta) = A_k \left[\alpha \frac{e^{-ik\eta}}{\sqrt{2k}} + \beta \frac{e^{ik\eta}}{\sqrt{2k}} \right], \quad (53)$$

where the coefficients α and β are determined by matching the solutions at the onset of the radiation stage.

We define “energy” of the harmonic oscillator as

$$\mathcal{E} := \frac{1}{2} |u'|^2 + \frac{k^2}{2} |u|^2. \quad (54)$$

Then, this energy is conserved at the zeroth order, i.e.,

$$\mathcal{E}^{(0)} = \frac{k}{2} (|\alpha|^2 + |\beta|^2) |A_k|^2 = \text{const.}$$

Taking into account the corrections of $\mathcal{O}(\ell^2)$, the total variation of the energy can be estimated as

$$\Delta \mathcal{E} = \sum_{j=0}^2 \Delta \mathcal{E}_j = \sum_{j=0}^2 \int_{\eta_0}^{\infty} \mathcal{E}'_j d\eta, \quad (55)$$

with

$$\begin{aligned} \mathcal{E}'_0 &= \frac{\ell^2}{2a} (u')^* \bar{\mathcal{S}}_0 + \text{c.c.}, \\ \mathcal{E}'_j &= \frac{\ell^2}{2a} (u')^* \mathcal{S}_j + \text{c.c.}, \quad \text{for } j = 1, 2. \end{aligned} \quad (56)$$

The amplitude of the oscillator $|A_k| e^{\Re[F]}$ is proportional to $\mathcal{E}^{1/2} a^{(0)}/a$. The factor $a^{(0)}/a$ here is to be attributed to unconventional cosmic expansion. Hence we have

$$\begin{aligned} \lim_{\eta \rightarrow \infty} \left\{ \Re[F_0(\eta)] + \frac{\delta a(\eta)}{a^{(0)}(\eta)} \right\} &= \frac{\Delta \mathcal{E}_0}{2\mathcal{E}^{(0)}}, \\ \lim_{\eta \rightarrow \infty} \Re[F_j(\eta)] &= \frac{\Delta \mathcal{E}_j}{2\mathcal{E}^{(0)}}, \quad \text{for } j = 1, 2. \end{aligned} \quad (57)$$

A. Corrections due to the unconventional expansion law

We start with computing the correction due to the unconventional cosmic expansion, F_0 , which we could not determine in the long wavelength approximation. To do so, we first need to determine the coefficients in Eq. (53) by requiring that $u^{(0)}$ and its derivative are continuous across the transition from the de Sitter stage to the radiation stage. We assume for a moment that the perturbation in the initial phase is given by the de Sitter growing mode,

$$v(\eta) \simeq \frac{A_k a_0 \eta_0}{2\eta_0 - \eta} \left[1 + \frac{k^2 (2\eta_0 - \eta)^2}{2} \right]. \quad (58)$$

Solving

$$\begin{aligned} v(\eta_0) &= {}^{(0)}u(\eta_0), \\ \partial_\eta v(\eta_0) &= \partial_\eta {}^{(0)}u(\eta_0), \end{aligned}$$

we obtain the coefficients as

$$\alpha \simeq \frac{ia_0}{\eta_0 \sqrt{2k}} \left[1 - \frac{2i}{3}(k\eta_0)^3 \right], \quad \beta = \alpha^*. \quad (59)$$

The energy of the harmonic oscillator at zeroth order is approximately given by

$$\mathcal{E}^{(0)} \simeq \frac{|A_k|^2 a_0^2}{2\eta_0^2}. \quad (60)$$

Next we integrate

$$\begin{aligned} \mathcal{E}'_0 &= \frac{i}{4} \frac{a''}{a} (|\alpha|^2 - |\beta|^2 - \alpha\beta^* e^{-2ik\eta} + \alpha^*\beta e^{2ik\eta}) |A_k|^2 + \text{c.c.} \\ &= \frac{a''}{a} \Im[\alpha\beta^* e^{-2ik\eta}] |A_k|^2 \end{aligned} \quad (61)$$

The scale factor correct up to $\mathcal{O}(\ell^2)$ is given in Eq. (6). Using this expression, we have

$$\frac{a''}{a} \simeq -\frac{\ell^2 H_0^2}{2} \frac{\eta_0^4}{\eta^6}. \quad (62)$$

Then, integrating Eq. (61), we have

$$\begin{aligned} \frac{\Delta \mathcal{E}_0}{2 \mathcal{E}^{(0)}} &\simeq -\frac{\ell^2 H_0^2 \eta_0^2}{2a_0^2} \Im \left[\alpha\beta^* \int_{\eta_0}^{\infty} d\eta \frac{\eta_0^4}{\eta^6} e^{-2ik\eta} \right] \\ &\simeq \frac{\ell^2 H_0^2}{4k\eta_0} \Im \left[\left(1 - \frac{2i}{3}(k\eta_0)^3 \right)^2 \left[\frac{1}{5} - \frac{i}{2}k\eta_0 - \frac{2}{3}(k\eta_0)^2 + \frac{2i}{3}(k\eta_0)^3 + \mathcal{O}((k\eta_0)^4) \right] \right] \\ &\simeq -\frac{1}{8} \ell^2 H_0^2 + \frac{1}{10} \frac{k^2 \ell^2}{a_0^2}. \end{aligned} \quad (63)$$

The first term in the last line is not suppressed for the $k \rightarrow 0$ limit. This is because this term arises due to the asymptotic behavior of $\delta a(\eta)$. In fact one can see from Eq. (57) that the expression for the physical amplitude of the perturbations e^{F_0} does not have this term, and finally we have

$$\lim_{\eta \rightarrow \infty} \Re[F_0(\eta)] = \frac{1}{10} \frac{k^2 \ell^2}{a_0^2}. \quad (64)$$

B. Suppression at sub-horizon scales

Here we shall show that corrections from the $E_{\mu\nu}$ term are suppressed at sub-horizon scales and hence the result obtained in the preceding section by using the long wavelength approximation suffices for our purpose.

Substituting $\phi = {}^{(0)}u / {}^{(0)}a$ into $\mathcal{S}_1[\phi]$, energy gain or loss due to this term is obtained as

$$\mathcal{E}'_1 = \frac{\ell^2}{2 {}^{(0)}a^2} \left[\frac{5}{\eta^3} |{}^{(0)}u'|^2 + \left(-\frac{5}{\eta^4} + \frac{k^2}{\eta^2} \right) ({}^{(0)}u')^* {}^{(0)}u \right] + \text{c.c.} \quad (65)$$

Note that $|{}^{(0)}u'|^2$ and $({}^{(0)}u')^* {}^{(0)}u$ consist of just constant parts and oscillating parts with constant amplitudes. Therefore, it is manifest that when we integrate $\mathcal{E}'_1$ the dominant contribution comes from around the lower boundary of the integral, $\eta \approx \eta_0$, and so corrections arising in the sub-horizon regime are suppressed compared to those imprinted beforehand in the super-horizon regime.

A similar expression for $\mathcal{E}'_2$ can be obtained as

$$\mathcal{E}'_2 = -\frac{\ell^2}{8} \frac{(u^{(0)})^*}{a} \int d\omega p^4 \tilde{\phi}^{(0)} e^{-i\omega\eta} \ln\left(\frac{p^2}{k^2}\right) + \frac{2\ell^2}{a^2} \left[\ln\left(\frac{k\ell}{2a^{(0)}}\right) + \gamma \right] \left[\frac{3}{\eta^3} |u^{(0)'}|^2 + \left(-\frac{3}{\eta^4} + \frac{k^2}{\eta^2}\right) (u^{(0)'})^* u^{(0)} \right] + \text{c.c.} \quad (66)$$

The second term is local and suppressed for large η for the same reason above. To examine the late time behavior of the first term, we substitute the expression (53) without limitation $\eta > \eta_0$. This approximation is justified because the kernel $\int d\omega e^{-i\omega(\eta-\eta')} p^4 \ln(p^2/k^2)$ decays at least as fast as $1/(\eta-\eta')^3$ for a large time separation, as will be shown in Appendix. Under this approximation for $\phi(\eta)$, it is easy to calculate its Fourier transform²

$$\tilde{\phi}^{(0)}(\omega) = \frac{i\eta_0}{2a_0} \frac{1}{\sqrt{2k}} \times \begin{cases} -(\alpha + \beta) & \text{for } \omega \leq -k, \\ (\alpha - \beta) & \text{for } |\omega| \leq k, \\ (\alpha + \beta) & \text{for } \omega \geq k. \end{cases} \quad (67)$$

The contour of the ω -integration in the first term of Eq. (66) is such shown in Fig. 3, and hence it can be separated into three parts as

$$\text{The first term in (66)} = -\frac{(u^{(0)})^*}{16} \frac{k^2 \ell^2}{a_0 a} k\eta_0 (\mathcal{I}_{\rightarrow -k}^{-k-i\infty} + \mathcal{I}_{\rightarrow k}^{-k} + \mathcal{I}_{\rightarrow k-i\infty}^k), \quad (68)$$

where

$$\mathcal{I}_{\rightarrow k-i\infty}^k = (\mathcal{I}_{\rightarrow -k}^{-k-i\infty})^* = (\alpha + \beta) \frac{e^{-ik\eta}}{\sqrt{2k}} \int_0^\infty dy (y^2 + 2iy)^2 e^{-k\eta \cdot y} \left[\ln(2y - iy^2) - \frac{\pi i}{2} \right], \quad (69)$$

$$\mathcal{I}_{\rightarrow k}^{-k} = (\alpha - \beta) \frac{i}{\sqrt{2k}} \int_{-1}^1 dy (1 - y^2)^2 e^{-ik\eta \cdot y} \ln(1 - y^2). \quad (70)$$

We can see that all $\mathcal{I}$ s are suppressed for large values of $k\eta$, since their integrand becomes a product of a sooth function and a rapidly oscillating function. The dominant contribution in the integration of $\mathcal{E}'$ therefore comes from around the lower boundary of the integral.

Based on the above analysis, we conclude that all the dominant corrections arise in the super-horizon regime, and thus it is sufficient to evaluate the corrections in the long wavelength approximation.

VI. SUMMARY AND DISCUSSION

We have investigated leading order corrections to the tensor perturbations in the RS II braneworld cosmology by using the perturbative expansion scheme of Ref. [28]. We have studied a model composed of slow-roll inflation on the brane, followed by a radiation dominant era. The unperturbed five dimensional bulk is AdS space with curvature radius ℓ . In our expansion scheme the asymptotic boundary conditions in the bulk are imposed by choosing outgoing solutions of bulk perturbations, whose general expression is known in the Poincaré coordinate system. Hence, the issue of bulk boundary conditions is handled without introducing an artificial regulator brane. This is one of the notable advantages of the present scheme.

We set the initial condition when the wave length ak^{-1} is already longer than the Hubble radius during slow-roll inflation. We will not lose much by neglecting the modes which are already inside the Hubble scale at the time of transition to the radiation era, since they are not cosmologically so interesting. We compared the resulting amplitudes of fluctuations after the horizon re-entry with and without the effect of an extra dimension. To do so, we normalized the amplitude so that the late time amplitude of fluctuations becomes identical in two cases if the slow-roll inflation lasts forever. As the reference without the effect of an extra dimension, we took a growing mode solution. The corrections due to gravity propagation through the fifth dimension dominantly come from the contribution around the transition time. Therefore they can be estimated by using the long wavelength approximation with a sufficient accuracy. Since the correction due to the modified cosmic expansion comes from relatively late epoch, it is necessary to take into account the oscillatory behavior of the solution after the re-entry to the Hubble horizon.

² Depending on the choice of the integration path near $\eta = 0$, the Fourier transform of $\phi(\eta)$ changes by a constant independent of ω . Here we took the principal values.

Combining the results obtained in Eqs. (43), (49) and (64), we find that the amplitude of a fluctuation with comoving wave number k is modified by a factor $e^{\Re[F]}$ due to the effect of an extra dimension with

$$\lim_{\eta \rightarrow \infty} \Re[F(\eta)] = \frac{k^2 \ell^2}{a_0^2} \left\{ \frac{13}{150} - \frac{4}{15} \left[\gamma + \ln \left(\frac{k}{a_0 H_0} \right) \right] + \frac{2}{5} \ln \left(\frac{\ell H_0}{2} \right) \right\}, \quad (71)$$

where a_0 and H_0 are the scale factor and the Hubble parameter at the transition time, and γ is the Euler's constant. Leading corrections are proportional to $k^2 \ell^2 / a_0^2$ or $k^2 \ell^2 / a_0^2 \log k$, as is expected from the dimensional analysis. However, our calculation here determined the precise numerical factors analytically.

Throughout this paper, we have dropped the contribution from the decaying mode for simplicity. Since the decaying mode during the initial slow-roll inflation phase is usually suppressed when the wavelength is longer than the Hubble scale, it does not give a significant effect. However, in the context of braneworld, we have not yet understood how to fix the initial conditions for fluctuations. Therefore, in principle, there is a possibility that the contamination of the decaying mode can be extremely large, although it seems quite unlikely. In such a case the contribution from the decaying component cannot be neglected.

It is quite easy to estimate the leading order correction due to the decaying mode, since it does not pick up the effect of the presence of an extra dimension. Here the effect is not due to the non-trivial evolution of a solution but totally due to unconventional initial condition. Adding a decaying mode will modify v given in Eq. (58) to

$$v(\eta) \simeq \frac{A_k a_0 \eta_0}{2\eta_0 - \eta} \left[1 + \frac{k^2 (2\eta_0 - \eta)^2}{2} + C_d k^3 (2\eta_0 - \eta)^3 \right],$$

where C_d is a complex number which parametrizes the amplitude of the decaying component. Repeating a similar calculation leading to Eq. (64), we find that the relative change in the amplitude due to the decaying mode is given by

$$\Re[C_d] \frac{\ell^2 H_0^2}{2} (k\eta_0)^3. \quad (72)$$

Thus the correction due to the decaying component is more suppressed in the sense of the power of k , but it might be more important than the other corrections for $k\eta_0 > \Re[C_d]^{-1}$ if $\Re[C_d]$ is extremely large.

Lastly we would like to mention the limitation of the present work and some future issues. The perturbative expansion scheme that we have adopted in this paper is valid for scales larger than the AdS curvature length ($k\ell/a \ll 1$) at low energies ($\ell H \ll 1$); besides our study is restricted to initially super-horizon perturbations. We gave an initial condition for a solution by hand, but of course such an approach is not satisfactory. To determine the initial condition, we need to investigate the evolution of perturbations at the initial phase where the wavelength of the mode is too short and reduction to an effective four dimensional problem is not possible. Furthermore, to seek for interesting effects that might arise at high energies ($\ell H \gtrsim 1$), we have to develop a new formalism, which is a next challenge and will be hopefully reported in our future publication.

Acknowledgments

We are grateful to Shinji Mukohyama and Roy Maartens for useful discussions. This work is supported in part by Monbukagakusho Grant-in-Aid Nos. 16740141 and 14047212, Inamori Foundation, and 21COE program at Kyoto university "Center for Diversity and Universality in Physics".

APPENDIX A: DETAILS OF CALCULATIONS

In this appendix, we will show the following formula:

$$\int d\omega e^{-i\omega T} \ln \left(\frac{p^2}{k^2} \right) = -4\pi \left\{ \theta(T) \frac{\cos(kT) - 1}{T} + \partial_T [\theta(T) \ln(kT)] + \gamma \delta(T) \right\}, \quad (A1)$$

where $p^2 = k^2 - \omega^2$ and γ is Euler's constant. The branch cuts are all running on the lower half complex plane of ω (Fig.3), which ensures the retarded boundary conditions.

We divide the above integration into two parts:

$$\int d\omega e^{-i\omega T} \ln \left(\frac{p^2}{k^2} \right) = \int d\omega e^{-i\omega T} \ln \left(\frac{-p^2}{\omega^2} \right) + \int d\omega e^{-i\omega T} \ln \left(\frac{-\omega^2}{k^2} \right). \quad (A2)$$

The first term of the right hand side can be rewritten as

$$\int d\omega e^{-i\omega T} \ln\left(\frac{-p^2}{\omega^2}\right) = \int d\omega e^{-i\omega T} [\ln(\omega - k) + \ln(\omega + k) - 2\ln\omega].$$

The integration can be performed as

$$\begin{aligned} \int d\omega e^{-i\omega T} \ln(\omega - k) &= \theta(T) \left[\int_{-\infty}^0 (-i) d\omega e^{-i\omega T} \ln(\omega - k) + \int_0^{\infty} (-i) d\omega e^{-i\omega T} \ln(\omega - k) \right] \\ &= -2\pi\theta(T) \frac{e^{-ikT}}{T}, \end{aligned}$$

and the other two integrals are obtained by setting $k \rightarrow -k$ and $k \rightarrow 0$. Thus we have

$$\int d\omega e^{-i\omega T} \ln\left(\frac{-p^2}{\omega^2}\right) = -4\pi\theta(T) \frac{\cos(kT) - 1}{T}.$$

This expression is regular at $T = 0$. To evaluate the second term in the right hand side of Eq. (A2), we need a trick because the integrand suffers from a slow fall-off at large values of $|\omega|$. It should be interpreted as

$$2i\partial_T \left[\int d\omega \frac{e^{-i\omega T}}{\omega} \ln\left(\frac{-i\omega}{k}\right) \right] = 2i\partial_T \left\{ \theta(T) \left[\int d\lambda \frac{e^{-\lambda}}{\lambda} (\ln(-\lambda) - \ln(kT)) \right] \right\}, \quad (\text{A3})$$

where we put $\lambda = i\omega T$. Now the branch cut and the contour of the integration on λ plane are such shown in Fig 4. Thanks to this trick, the expression for the second term becomes a well-defined one, though the result does not change as long as convergence is guaranteed. Then, performing the integration as

$$\begin{aligned} \int d\lambda \frac{e^{-\lambda}}{\lambda} (\ln(-\lambda) - \ln(kT)) &= \lim_{\nu \rightarrow 0} \partial_{\nu} \int d\lambda e^{-\lambda} (-\lambda)^{\nu-1} + 2\pi i \ln(kT) \\ &= \lim_{\nu \rightarrow 0} \partial_{\nu} [(e^{-\pi i \nu} - e^{\pi i \nu}) \Gamma(\nu)] + 2\pi i \ln(kT) \\ &= 2\pi i [\ln(kT) + \gamma], \end{aligned}$$

we finally obtain the formula (A1).

For $T > 0$ the formula (A1) reduces to $\int d\omega e^{-i\omega T} \ln(p^2/k^2) = -4\pi \cos(kT)/T$ and hence the kernel that appears in $\mathcal{S}_2$ becomes

$$\int d\omega e^{-i\omega T} p^4 \ln\left(\frac{p^2}{k^2}\right) = 32\pi \left[\left(\frac{k^2}{T^3} - \frac{3}{T^5} \right) \cos(kT) - \frac{3k}{T^4} \sin(kT) \right], \quad \text{for } T > 0. \quad (\text{A4})$$

From this expression we find that the non-local source $\mathcal{S}_2$ does not keep the past history for a long time.




FIG. 4: Branch cut and the contour of the integration in Eq. (A3).

- [2] D. Langlois, Prog. Theor. Phys. Suppl. **148**, 181 (2003) [arXiv:hep-th/0209261].
- [3] R. Maartens, “Brane-world gravity,” arXiv:gr-qc/0312059.
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
- [5] J. Garriga and T. Tanaka, Phys. Rev. Lett. **84**, 2778 (2000) [arXiv:hep-th/9911055].
- [6] S. B. Giddings, E. Katz and L. Randall, JHEP **0003**, 023 (2000) [arXiv:hep-th/0002091].
- [7] R. Maartens, D. Wands, B. A. Bassett and I. Heard, Phys. Rev. D **62**, 041301 (2000) [arXiv:hep-ph/9912464].
- [8] S. Mukohyama, Phys. Rev. D **62**, 084015 (2000) [arXiv:hep-th/0004067].
- [9] H. Kodama, A. Ishibashi and O. Seto, Phys. Rev. D **62**, 064022 (2000) [arXiv:hep-th/0004160].
- [10] D. Langlois, Phys. Rev. D **62**, 126012 (2000) [arXiv:hep-th/0005025].
- [11] D. Langlois, Phys. Rev. Lett. **86**, 2212 (2001) [arXiv:hep-th/0010063].
- [12] K. Koyama and J. Soda, Phys. Rev. D **62**, 123502 (2000) [arXiv:hep-th/0005239].
- [13] K. Koyama and J. Soda, Phys. Rev. D **65**, 023514 (2002) [arXiv:hep-th/0108003].
- [14] K. Koyama, Phys. Rev. Lett. **91**, 221301 (2003) [arXiv:astro-ph/0303108].
- [15] D. Langlois, R. Maartens and D. Wands, Phys. Lett. B **489**, 259 (2000) [arXiv:hep-th/0006007].
- [16] D. S. Gorbunov, V. A. Rubakov and S. M. Sibiryakov, JHEP **0110**, 015 (2001) [arXiv:hep-th/0108017].
- [17] T. Kobayashi, H. Kudoh and T. Tanaka, Phys. Rev. D **68**, 044025 (2003) [arXiv:gr-qc/0305006].
- [18] T. Hiramatsu, K. Koyama and A. Taruya, Phys. Lett. B **578**, 269 (2004) [arXiv:hep-th/0308072].
- [19] R. Easther, D. Langlois, R. Maartens and D. Wands, JCAP **0310**, 014 (2003) [arXiv:hep-th/0308078].
- [20] R. A. Battye, C. Van de Bruck and A. Mennim, Phys. Rev. D **69**, 064040 (2004) [arXiv:hep-th/0308134].
- [21] K. Ichiki and K. Nakamura, “Causal structure and gravitational waves in brane world cosmology,” arXiv:hep-th/0310282.
- [22] K. Ichiki and K. Nakamura, “Stochastic gravitational wave background in brane world cosmology,” arXiv:astro-ph/0406606.
- [23] K. Koyama, “Late time behavior of cosmological perturbations in a single brane model,” arXiv:astro-ph/0407263.
- [24] K. Koyama and K. Takahashi, Phys. Rev. D **67**, 103503 (2003) [arXiv:hep-th/0301165].
- [25] K. Koyama and K. Takahashi, Phys. Rev. D **68**, 103512 (2003) [arXiv:hep-th/0307073].
- [26] T. Kobayashi and T. Tanaka, Phys. Rev. D **69**, 064037 (2004) [arXiv:hep-th/0311197].
- [27] P. Binetruy, M. Bucher and C. Carvalho, “Models for the brane-bulk interaction: Toward understanding braneworld” arXiv:hep-th/0403154.
- [28] T. Tanaka, “AdS/CFT correspondence in a Friedmann-Lemaitre-Robertson-Walker brane,” arXiv:gr-qc/0402068.
- [29] P. Binetruy, C. Deffayet and D. Langlois, Nucl. Phys. B **565**, 269 (2000) [arXiv:hep-th/9905012].
- [30] P. Binetruy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. B **477**, 285 (2000) [arXiv:hep-th/9910219].
- [31] S. Mukohyama, Phys. Lett. B **473**, 241 (2000) [arXiv:hep-th/9911165].
- [32] D. Ida, JHEP **0009**, 014 (2000) [arXiv:gr-qc/9912002].
- [33] P. Kraus, JHEP **9912**, 011 (1999) [arXiv:hep-th/9910149].
- [34] T. Shiromizu, K. i. Maeda and M. Sasaki, Phys. Rev. D **62**, 024012 (2000) [arXiv:gr-qc/9910076].
- [35] S. Kanno and J. Soda, Phys. Rev. D **66**, 043526 (2002) [arXiv:hep-th/0205188].
- [36] S. Kanno and J. Soda, Phys. Rev. D **66**, 083506 (2002) [arXiv:hep-th/0207029].
- [37] K. Koyama, Phys. Rev. D **66**, 084003 (2002) [arXiv:gr-qc/0204047].